

Hints of a Local Matter Underdensity or Modified Gravity in the Low z Pantheon data

L. Kazantzidis* and L. Perivolaropoulos†

Department of Physics, University of Ioannina, GR-45110, Ioannina, Greece

(Dated: April 7, 2020)

A redshift tomography of the Pantheon type Ia supernovae (SnIa) data focusing on the best fit value of the absolute magnitude M and/or Hubble constant H_0 in the context of Λ CDM indicates a local variation ($z \lesssim 0.2$) at 2σ level, with respect to the best fit of the full dataset. If this variation is not due to a statistical fluctuation, it can be interpreted as a locally higher value of H_0 by about 2%, corresponding to a local matter underdensity $\delta\rho_0/\rho_0 \simeq -0.10 \pm 0.04$. It can also be interpreted as a time variation of Newton's constant which implies an evolving Chandrasekhar mass and thus an evolving absolute luminosity L and absolute magnitude M of low z SnIa. The local void scenario would predict a degree of anisotropy in the best fit value of H_0 (and/or M) since it is unlikely that we are located at the center of a local spherical underdensity. Using a hemisphere comparison method, we find an anisotropy level that is consistent with simulated isotropic Pantheon-like datasets. We show however, that the anisotropic sky distribution of the Pantheon SnIa data induces a preferred range of directions even in simulated Pantheon data obtained in the context of isotropic Λ CDM. We thus construct a more isotropically distributed subset of the Pantheon SnIa and show that the preferred range of directions disappears. Using this more isotropically distributed subset we again find no evidence for statistically significant anisotropy using either the hemisphere comparison method or the dipole fit method. In the context of the modified gravity scenario, we allow for an evolving normalized Newton's constant consistent with General Relativity (GR) at early and late times $\mu(z) = G_{\text{eff}}(z, g_a)/G_N = 1 + g_a z^2/(1+z)^2 - g_a z^4/(1+z)^4$ and fit for the parameter g_a assuming $L \sim G_{\text{eff}}^b$. For $b = -3/2$ indicated by some previous studies we find $g_a = -0.47 \pm 0.36$ which is more than 1.5σ away from the GR value of $g_a = 0$. This weak hint for weaker gravity at low z coming from SnIa is consistent with similar evidence from growth and weak lensing cosmological data.

I. INTRODUCTION

Since the discovery of the accelerating expansion of the universe [1, 2], the Λ CDM model based on the existence of a cosmological constant [3] has been particularly simple and consistent with most cosmological observations including Cosmic Microwave Background (CMB) perturbations [4–6], Type Ia supernovae (SnIa) standard candles [7, 8] and Cosmic Chronometer probes of the expansion rate $H(z)$ [9–11], Baryon Acoustic Oscillations (BAO) standard ruler probes of $H(z)$ [12, 13], Large Scale Matter perturbations observed through Redshift Space Distortions (RSD) [6, 14, 15], Weak Lensing (WL) [16, 17], Cluster Count data [18–20] etc. Despite of its overall success and simplicity, the Λ CDM model faces challenges on both theoretical and observational grounds. Theoretical challenges of Λ CDM include the fine tuning [21, 22] and coincidence [23, 24] problems. Observationally, there have been indications that different cosmological observations favour different values for the basic parameters of the model (at a level of 2σ or more) [25–28] indicating that new degrees of freedom may be required to make the model simultaneously consistent with all these observations. These “tensions” of Λ CDM include the following:

- **The H_0 problem $> 4\sigma$:** CMB and BAO cosmological measurements using the last scattering sound horizon as a standard ruler and assuming a Λ CDM background expansion, report $H_0 = 67.4 \pm 0.5 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [6], a best fit value which is about 9% lower compared to the local measurement of H_0 coming from SnIa data, that publish $H_0 = 74.03 \pm 1.42 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [29]. The discrepancy ranges from 4.4σ to more than 5σ [30, 31] depending on the combination of local data considered. Nevertheless, independent measurements of cosmic chronometers (based on models of evolving galaxy star luminosity) report a best fit value of $H_0 = 67.06 \pm 1.68 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [11] favouring the CMB and BAO measurements. On the contrary, measurements of H_0 based on a calibration of the Tip of the Red Giant Branch which is applied on SnIa (instead of the Cepheid calibration method), report $H_0 = 69.8 \pm 0.8 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [32], a value that differs approximately 3σ away from the best fit values that describe the H_0 problem.
- **The growth tension $\simeq 3\sigma$:** The growth rate and magnitude of linear cosmological perturbations depend on the matter density parameter Ω_{0m} and on the amplitude of the primordial power spectrum which is measured through the parameter σ_8 , the linear amplitude of matter fluctuations on scales $8h^{-1} \text{ Mpc}$. Weak Lensing (WL) [33–37] and Redshift Space Distortion (RSD) [38–47] cosmological observations measuring directly the growth rate of

* l.kazantzidis@uoi.gr

† leandros@uoi.gr

cosmological perturbations (dynamical probes) indicate that the observed growth rate is weaker than expected in the context of Λ CDM with parameters determined from the observed background expansion rate using geometric probes (SnIa, BAO and CMB standard ruler data). This discrepancy is expressed as a preference for lower values of the parameters Ω_{0m} and σ_8 by dynamical probes compared to the corresponding values favoured by the geometric probes. The level of the growth tension is at least about $2 - 3\sigma$ [38–40, 43–45] but it can vary up to about 5σ (when the E_G statistic data are used [48]) depending on the model parametrization and the type of dataset considered.

- **Low- z galaxy BAO vs high z Ly α BAO curvature ($\simeq 2\sigma$):** There is a $\sim 2\sigma$ tension [49, 50] between the value of Ω_{0m} favoured by Ly- α BAO measurements ($\Omega_{0m} \simeq 0.19 \pm 0.07$ for $z > 2.4$), which favour lower values of Ω_{0m} , and the values of Ω_{0m} favoured by galaxy BAO measurements ($\Omega_{0m} \simeq 0.37 \pm 0.07$ for $z < 0.6$) that favour higher values of Ω_{0m} .
- **Low l – high l CMB power spectrum curvature ($\simeq 2\sigma$):** There is a mismatch of the cold dark matter density parameter $\Omega_c h^2$ best fit that is derived using high ($l > 1000$) and low multipoles ($l < 1000$). This tension is approximately at a 2.5σ level [51, 52] and is such that the low l multipoles predict a lower value of $\Omega_c h^2$ than the high- l multipoles. It is also described by the need to introduce the A_L parameter [52] which multiplies the amplitude of the lensing potential power [52]. Thus, this tension is also described by the fact that the high- l TT multipoles are observed to correspond to a higher $\phi\phi$ lensing potential (the high- l secondary peaks of the TT CMB power spectrum are smoother than expected in the context of the best fit Planck15/ Λ CDM model parameters). The value of the best fit Hubble parameter is also about 2.5σ lower when obtained from the higher- l multipoles ($H_0 = 64.1 \pm 1.7 \text{ km s}^{-1} \text{ Mpc}^{-1}$) compared to the corresponding best fit value obtained from the low- l CMB spectrum multipoles ($H_0 = 69.7 \pm 1.7 \text{ km s}^{-1} \text{ Mpc}^{-1}$) [51].

The strongest of the above tensions which has also been called a “problem” due its persistence in time and its increasing statistical significance is the Hubble parameter tension. This is heavily based on the use of SnIa as standard candle probes of the cosmic expansion rate. SnIa have been extensively used as standard candles to probe the expansion rate (Hubble parameter) $H(z)$ of the late universe ($z < 2$). The theoretically predicted apparent magnitude $m_{th}(z)$ of SnIa is connected with the Hubble

free luminosity distance $D_L(z) \equiv H_0 d_L(z)/c$ as

$$m_{th}(z) = M + 5 \log_{10} [D_L(z)] + 5 \log_{10} \left(\frac{c/H_0}{1 \text{ Mpc}} \right) + 25 \quad (1.1)$$

where M is the colour and stretch corrected absolute magnitude of SnIa (assumed constant) and $d_L(z)$ is the luminosity distance of each SnIa which in a flat universe is

$$d_L(z) = c(1+z) \int_0^z \frac{dz'}{H(z')} \quad (1.2)$$

Using Eq. (1.1), measurements of the SnIa apparent magnitude $m(z)$ at various redshifts can be used to determine the present day Hubble parameter H_0 as well as its redshift dependence through Eq. (1.2). For the determination of H_0 , Riess et. al [53] used local distance ladder measurements (Cepheid calibrations at $z \simeq 0.01$) to measure directly M and then a kinematic local expansion of $D_L(z)$ as

$$D_L(z) = z \left[1 + \frac{1}{2}(1 - q_0)z - \frac{1}{6}(1 - q_0 - 3q_0^2 + j_0)z^2 + \dots \right] \quad (1.3)$$

to fit for the parameters H_0 , q_0 , j_0 [54] using low z SnIa ($z \lesssim 0.2$).

For the determination of cosmological parameters in $H(z)$, higher z SnIa are used and the degenerate parameters M , H_0 are usually marginalized as nuisance parameters [7, 8, 55]. For example, in the context of Λ CDM with

$$H^2(z) = H_0^2 [\Omega_{0m}(1+z)^3 + (1 - \Omega_{0m})] \quad (1.4)$$

Eq. (1.1) is used for the construction and minimization of $\bar{\chi}^2(\Omega_{0m}) \equiv \int d\mathcal{M} \chi^2(\mathcal{M}, \Omega_{0m})$ where the degenerate combination

$$\mathcal{M} \equiv M + 5 \log_{10} \left[\frac{c/H_0}{1 \text{ Mpc}} \right] + 25 = M - 5 \log_{10}(h) + 42.38 \quad (1.5)$$

($H_0 = 100h \text{ km s}^{-1} \text{ Mpc}^{-1}$) has been marginalized.

The marginalization of the parameter \mathcal{M} however can lead to loss of useful physical information related to possible spatial variations of H_0 and/or time variations of the absolute magnitude M . For example, a value of \mathcal{M} that evolves with redshift in a way that leads to low \mathcal{M} values at low z could imply either higher local values of H_0 due to a local matter underdensity or lower values of the absolute magnitude M at recent cosmological times due to *e.g.* a time variation of Newton’s constant.

The former case is in agreement with a few independent groups that have found evidence for a local matter underdensity on scales $100 - 300h^{-1} \text{ Mpc}$ [56, 57] with $\delta\rho_0/\rho_0$ in the range between -0.1 and -0.3 using either SnIa [58] in the context of a Lemaitre-Tolman Bondi (LTB) metric [59–61] ($\delta\rho_0/\rho_0 \simeq -0.15$) or galaxy survey catalogues [62] to construct luminosity density samples in the redshift range of $0.01 < z < 0.2$ ($\delta\rho_0/\rho_0 \simeq -0.3$).

A local matter underdensity of about 15% corresponds to a local variation (increase) of H_0 by about 2% which is in the right direction but not large enough to explain the H_0 tension which would require a local increase of H_0 by about 9% compared to its mean value in the universe, *i.e.* a much deeper underdensity than the one implied by SnIa data.

If the later case is realized in Nature, the evolution of the absolute magnitude M of SnIa (or equivalently the absolute luminosity $L \sim 10^{-2M/5}$) could be used as a probe of the evolution of fundamental constants like the fine structure α or the Newton's constant G_{eff} . In the physical context of an evolving G_{eff} , previous studies [63, 64] assumed that the amount of ${}^{56}\text{Ni}$ that is produced in a SnIa and determines the absolute luminosity L , is proportional to the Chandrasekhar mass $m_{ch} \sim G_{\text{eff}}^{-3/2}$ which implies that L will increase as G_{eff} decreases. In contrast, other more recent studies [65] using a semi-analytical model to obtain SnIa light curves in the context of modified gravity, have indicated that L will increase as G_{eff} increases. Assuming a power law dependence $L(z) \sim G_{\text{eff}}(z)^b$ and fixing the value of b , any detected redshift dependence of SnIa absolute luminosity (or equivalently absolute magnitude) can be translated into a redshift dependence of G_{eff} .

Therefore, a possible detection of redshift dependence of the parameter \mathcal{M} could imply either a local underdensity and spatially varying H_0 or a redshift dependent M and thus, possibly, an evolving G_{eff} . Since M and H_0 are degenerate parameters within \mathcal{M} in Eq. (1.5), the two scenarios can not be distinguished using only the redshift dependence of the SnIa apparent magnitudes. However, the local matter underdensity scenario with an off-center observer generically predicts a level of anisotropy in the best fit value of the parameter \mathcal{M} which would emerge due to the anisotropy of H_0 which is expected for an off center observer in a region of matter underdensity. Such an anisotropy could also manifest itself as an anisotropy of cosmological parameters entering $H(z)$ like the matter density parameter Ω_{0m} . Despite intense efforts to identify such anisotropy in the latest SnIa data (the Pantheon compilation [66–70] and the joint light-curve analysis of the SDSS-II and SNLS supernova samples data (JLA) [71–73]) no such anisotropy has been identified at a statistically significant level [66–73]. However, we stress that none of these analyses has attempted to identify anisotropy signals using the parameter \mathcal{M} (or equivalently the parameters H_0 and/or M). In the present analysis we aim to fill this gap in the literature¹.

The main questions addressed in the present analysis include the following:

- What is the level of statistical significance for an evolving with redshift parameter \mathcal{M} in the context of the Pantheon SnIa dataset?

- Are there any hints for anisotropy for the parameter \mathcal{M} (or equivalently the parameters M and/or H_0) in the context of the Pantheon SnIa dataset? Such an anisotropy would favour the scenario of a local matter underdensity rather than evolving absolute magnitude M . What is the optimal method for detecting such a possible anisotropy?
- If any hint for evolving \mathcal{M} is interpreted as a hint for evolving M and evolving Newton constant what is the best fit of the evolving $\mu(z) \equiv G_{\text{eff}}(z)/G_{\text{N}}$ (where G_{N} is the value of Newton's constant measured on solar system scales) and does it correspond to weakening gravity at low z as the growth and weak lensing cosmological data seem to indicate?

The structure of this paper is the following: In the next section we use various subsets of the Pantheon dataset, to obtain the possible redshift dependence of the best fit parameters \mathcal{M} and Ω_{0m} in the context of Λ CDM. In section III we use the hemisphere comparison (HC) method and the dipole fitting (DF) method to search for possible statistically significant directional dependence of the best fit parameter \mathcal{M} (or equivalently the parameter H_0 with fixed M). Such an anisotropy would be generically anticipated in the context of a cosmological off-center observer in a local matter underdensity. In section IV we make the assumption that any variation of the parameter \mathcal{M} is due to a variation of M induced by a varying $\mu(z)$ and in the context of a physically motivated single parameter parametrization of $\mu(z)$ we identify the best fit parameter value and corresponding strength of gravity at low z ($z \lesssim 0.2$) compared to the corresponding value at higher z . Finally, in section V, we summarize, discuss the possible physical implications of our results and identify the possible extensions of this work.

II. SEARCHING FOR A REDSHIFT DEPENDENCE OF \mathcal{M}

The Pantheon dataset [8] is the largest compilation to date that incorporates data from six different probes giving a total of 1048 SnIa datapoints covering the redshift range $0.01 < z < 2.3$. The publicly available data include the name of each SnIa, the redshifts in the CMB and heliocentric frames as well as the observed corrected apparent magnitude m_{obs} along with the corresponding error $\sigma_{m_{\text{obs}}}$. The m_{obs} of each SnIa is reported after applying color and stretch corrections as well as corrections due to biases from simulations of the SnIa. In the context of a maximum likelihood analysis [74] Eqs. (1.1), (1.2) and (1.5) are used to construct the appropriate χ^2 function as

$$\chi^2(\mathcal{M}, \Omega_{0m}) = V_{\text{Panth.}}^i C_{ij}^{-1} V_{\text{Panth.}}^j \quad (2.1)$$

where $V_{\text{Panth.}}^i \equiv m_{\text{obs}}(z_i) - m_{\text{th}}(z)$ and C_{ij} is the covariance matrix which is given as $C_{ij} = \bar{D}_{ij} + \bar{C}_{\text{sys}}$, where

¹ Notice however that in the case of the DF method, the parameter \mathcal{M} was taken into account in Refs. [69, 70]




FIG. 1: The evolution of the best fit values (blue dots) of \mathcal{M} (left panel), M (middle panel) and h (right panel) along with its 1σ error for various cutoff values z_{max} . The dashed lines correspond to the best fit values indicated by the full dataset.

\bar{D}_{ij} is the diagonal matrix

$$\bar{D}_{ij} = \begin{pmatrix} \sigma_{m_{obs,1}}^2 & 0 & 0 & \cdots \\ 0 & \sigma_{m_{obs,2}}^2 & 0 & \cdots \\ 0 & 0 & \cdots & \sigma_{m_{obs,N}}^2 \end{pmatrix} \quad (2.2)$$

and \bar{C}_{sys} is a non-diagonal matrix associated with the systematic uncertainties that emerge from the bias corrections method (see Ref. [8] for more details). The magnitude of the \bar{C}_{sys} elements is much smaller than the elements of \bar{D}_{ij} and thus we ignore it in the present analysis for simplicity. We have shown however that the effect of this approximation on our results is negligible.

As discussed in the Introduction there is a power law dependence of the absolute luminosity L on G_{eff} leading to a simple power law relation between M and G_{eff} . For $L \sim G_{eff}^b$ this equation is of the form

$$M - M_0 = -\frac{5b}{2} \log_{10}(\mu) \quad (2.3)$$

where M_0 corresponds to a reference local value of the absolute magnitude. Then, Eq. (1.1) takes the following form

$$m_{th}(z) = \mathcal{M} + 5 \log_{10}[D_L(z)] - \frac{5b}{2} \log_{10}(\mu) \quad (2.4)$$

where \mathcal{M} is given in Eq. (1.5) with M replaced by M_0 .

Most previous studies used $b = -3/2$ [63, 64] based on the assumption that $L \sim m_{ch} \sim G_{eff}^{-3/2}$. However, as mentioned in the Introduction, a more detailed analysis has been performed in [65], where the authors studied the effects of modified theories of gravity to the absolute magnitude M of the SnIa. In particular, in the semi-analytic model that was used, extra parameters such as the initial nickel mass in the ejecta, the initial radius of shock breakout, the scale velocity, the effective opacity as well as total ejecta mass were included. Then, the generated light curves were standardised by rescaling the shape to match a template width and the numerical dependence of the standardised intrinsic absolute luminosity L on G_{eff} was identified. Using this semi-analytical method a new power law relation between L and G_{eff} was derived with $b > 0$ (see the left panel of Fig. 7 of [65]).

A marginalisation is usually implemented over \mathcal{M} in most analyses of the SnIa data (e.g. [55]). This approach however, may lead to loss of useful information regarding possible redshift dependence of H_0 and/or M and G_{eff} . Thus, we choose to keep this parameter and fit it along with the cosmological parameter Ω_{0m} . Fixing the background to that of a Λ CDM, we implement the maximum likelihood method [74] to obtain the best fit values for Ω_{0m} and \mathcal{M} as $\mathcal{M} = 23.803 \pm 0.007$ and $\Omega_{0m} = 0.285 \pm 0.012$ for the full Pantheon dataset in agreement with previous studies [8, 69].

In the context of a redshift independent \mathcal{M} and a Λ CDM background, any subset of the Pantheon dataset should provide best fit parameter values for Ω_{0m} and \mathcal{M} consistent with the corresponding best fit values of the full dataset. In order to test this conjecture, we fix Ω_{0m} to its best fit value indicated by the full dataset and consider subsets of the full dataset in redshift ranges $z \in [z_{min}, z_{max}]$ where $z_{min} = 0.02$ (fixed) and $z_{max} \geq 0.03$ (increasing for each point in steps of $\Delta z_{max} = 0.01$) is a cutoff redshift chosen so that the subsamples have acceptable statistics (the first and smallest subsample with $z_{max} = 0.03$ has 46 datapoints). Using each subsample, we implement the maximum likelihood method to find the best fit \mathcal{M} values along with their 1σ errors shown in Fig. 1 (left panel). For the best fit values of M of each subsample (Fig. 1 middle panel) we fix $h = 0.74$, i.e. to the value specified in [29], while for the best fit values of h (right panel of Fig. 1) we fix M using the best fit value of \mathcal{M} indicated by the full dataset and $h = 0.74$.

Clearly, at low redshifts and in particular in the redshift range $z_{max} \in [0.02, 0.15]$ there is a tension of about 2σ or more between the best fit value of \mathcal{M} of each subsample and the best fit value indicated by the full dataset. This difference may imply a lower value of M (middle panel of Fig. 1) or equivalently a higher value of h (right panel of Fig. 1) in the same redshift range. For $z_{max} > 0.15$ the best fit values in each subsample are consistent with the values indicated by the full dataset (dashed lines in Fig. 1) within 1σ level.

A similar behaviour is detected, if we rank the SnIa data from lowest to highest redshifts. At first we select the first 100 datapoints and fixing the background to the




FIG. 2: The evolution of the best fit values (blue dots) of \mathcal{M} (left panel), M (middle panel) and h (right panel) along with its 1σ error for 100 point subsamples vs the mean redshift z_{mean} . The dashed lines correspond to the best fit values indicated by the full dataset.




FIG. 3: The best fit values of \mathcal{M} (left panel) and Ω_{0m} (right panel) as well as the 1σ errors for the four bins. The horizontal axis corresponds to the redshift range of each bin. The dashed line describes the best fit value of the full dataset while the dot dashed lines its 1σ error.

best fit Λ CDM $H(z)$, [Eq. (1.4) with $\Omega_{0m} = 0.285$], we find the best fit value of \mathcal{M} along with its 1σ error for the lowest redshift subsample. Then, we shift the 100 points subsample by one datapoint towards higher redshifts to produce the next point and continue until we cover the entire redshift range of the Pantheon dataset (Fig. 2 - left panel). The redshift z_{mean} shown in the horizontal axis, corresponds to the mean redshift value of each of the 100 point subsamples.

From Fig. 2 we observe that for $z_{mean} < 0.3$, the best fit value of \mathcal{M} oscillates around the best fit value of the full dataset at a level of about $1 - 2\sigma$ which may indicate a similar oscillating behaviour for M (middle panel) and/or h (right panel) in the same redshift range. In this case, the redshift range of the oscillation is larger than the redshift variation detected in Fig. 1, because as the cutoff redshift increases, so does the size of the corresponding subsample, leading to a cancellation of the oscillating effect in Fig. 1.

In order to improve the statistics of the low z subsamples and further investigate the observed tension at low z , we sort the Pantheon data from lowest to highest redshift and divide them in four equal uncorrelated bins consisting of 262 datapoints. Then, we apply the maximum likelihood method in each bin separately, considering a

Λ CDM background and leaving the parameters \mathcal{M} and Ω_{0m} to vary simultaneously. Minimizing Eq. (2.1), we derive the best fit values of Ω_{0m} and \mathcal{M} as well as the corresponding 1σ error for each bin as it is shown in Fig. 3.

Clearly a similar oscillating behaviour for \mathcal{M} is apparent as in Fig. 2. Furthermore, the best fit values of Ω_{0m} and \mathcal{M} derived from the lowest z bin ($0.01 < z < 0.13$) are more than 2σ lower than the best fit values indicated by the full dataset, in agreement with Figs. 1 and 2. This is also evident in Fig. 4, where the $1\sigma - 3\sigma$ contours of the four bins are constructed in the parametric space ($\Omega_{0m} - \mathcal{M}$).

From Fig. 3 we find the difference of \mathcal{M} to be

$$\Delta\mathcal{M} \equiv \mathcal{M}_{bf} - \mathcal{M}_{bin1} \approx 23.80 - 23.76 \approx 0.04 \pm 0.02 \quad (2.5)$$

where \mathcal{M}_{bf} corresponds to the best fit value of \mathcal{M} indicated by the full dataset and \mathcal{M}_{bin1} corresponds to the best fit value of \mathcal{M} derived from the lowest z bin. In the context of a local matter underdensity, \mathcal{M}_{bf} is the true global value of \mathcal{M} , while \mathcal{M}_{bin1} corresponds to the value of \mathcal{M} that is measured in the interior of the local underdensity. This difference can be associated with a




FIG. 4: The $1\sigma - 3\sigma$ confidence contours in the parametric space $(\Omega_{0m} - \mathcal{M})$. The blue contours correspond to the $1\sigma - 3\sigma$ full Pantheon dataset best fit, while the red contours describe the $1\sigma - 3\sigma$ confidence contours of the four bins (from left to right). The black points represent the best fit of each bin, while the green dot represents the best fit value indicated by the full Pantheon dataset ($\Omega_{0m} = 0.285$ and $\mathcal{M} = 23.803$).




FIG. 5: The distributions of Ω_{0m} (left panel) and \mathcal{M} (right panel) using Monte Carlo simulations of Pantheon-like datasets under the assumption of an underlying Λ CDM model, in the redshift region $0.01 < z < 0.13$ (first bin). The red dashed lines correspond to the best fit values of the first bin.

variation of the local expansion rate $\delta H_0/H_0$ through

$$\left(\frac{\delta H_0}{H_0}\right) \approx 0.2 \ln(10) \Delta \mathcal{M} = 0.019 \pm 0.007 \quad (2.6)$$

We can also estimate the probability that such low values of \mathcal{M} and Ω_{0m} would occur in the context of the Λ CDM standard model, using Monte Carlo simulations of Pantheon-like datasets under the assumption of an underlying Λ CDM model with \mathcal{M} and Ω_{0m} corresponding to the values indicated by the full dataset. In particular, we construct 500 simulated datasets with redshifts corresponding to the redshifts of the first bin and substitute the apparent magnitude of the real data $[m_{obs}(z_i)]$ with simulated datapoints $m_{sim}(z_i)$ obtained from a random normal distribution with a mean value obtained from the best fit Λ CDM value of the apparent magnitude m_{th} [setting $\mathcal{M} = 23.803$ and $\Omega_{0m} = 0.285$ in Eq. (1.1)]. The standard deviation of the normal distribution is obtained from the $\sigma_{m_{obs}}$ of each datapoint respectively [75]. Then, we apply the maximum likelihood method and count how

many of the simulated data give lower values for \mathcal{M} and Ω_{0m} than the best fit values indicated by the real data of the first bin (red dashed lines). The results are plotted in Fig. 5. Clearly, less than 1% (0.2% for either Ω_{0m} or \mathcal{M}) of the Monte Carlo data give smaller best fit values for \mathcal{M} or Ω_{0m} than the actual best fit values of the first bin. Therefore, we confirm that this reduced value of \mathcal{M} is a highly unlikely event in the context of an underlying physical Λ CDM model.

The $2 - 3\sigma$ effect regarding the parameters Ω_{0m} and \mathcal{M} observed at low z have been also discussed in previous studies [58, 76]. In particular, for Ω_{0m} a similar behaviour was presented in [76], where the best fit values of Ω_{0m} and H_0 were studied, for various redshift cutoffs. Similar results for \mathcal{M} were also presented in [58], where the authors divided the Pantheon dataset in three bins and calculated the best fit value of $\mathcal{M} - 25$ in the context of a LTB model with a cosmological constant, in an attempt to identify hints of a local underdensity using the Pantheon dataset.

This variation of \mathcal{M} at low redshifts could be due to following:

- Some unknown statistical/systematic effect in the Pantheon dataset which affects the results at low z and has so far escaped attention.
- A local underdensity dubbed “Local Void” that fades away at large scales. Since \mathcal{M} is lower than the best fit value indicated by the full dataset at low z , h would be larger than the best fit value of the full dataset in the same redshift range. A generic way to explain this increase of h would be if our neighbourhood is more underdense compared to the mean density of the universe and as a result the measured value of h is also affected at local scales. In the context of a “Local Void” model, the value of h increases by 2 – 3% (see Fig. 2 - right panel). Such a scenario would also predict an anisotropy for the best fit value of \mathcal{M} in the sky.
- A modified theory of gravity. Another possible explanation for the observed variation of \mathcal{M} at low redshifts is a redshift dependence of M which could be due to a time variation of Newton’s constant in the context of a modified theory of gravity.

These two possibilities will be discussed in the next two sections.

III. “LOCAL VOID” SCENARIO

The idea that we live in an underdense region that fades away at large scales is not new. In fact, it has been proposed as an alternative theory to explain the accelerated expansion of the universe without the presence of a cosmological constant (see *e.g.* Refs. [77–81]). It has been shown however, the Gpc scale and depth of the underdensity required to explain the observed accelerating expansion is inconsistent with current observations [82]. Nevertheless, over the past twenty years, some works using various galaxy survey catalogues (*e.g.* the 2MASS survey [83, 84], the UKIDSS-Large Area Survey [85] as well as galaxies samples constructed from the 6dFGS, SDSS and GAMA surveys [86, 87]) have found some evidence for the existence of a local underdensity that extends on scales 150 – 400 $h^{-1}Mpc$ with depth $-0.4 < \delta\rho_0/\rho_0 < -0.05$. Other works considering the Pantheon dataset [58] or a sample of 1653 X-ray galaxy clusters [62] also stressed that a local underdensity on scales of $\approx 100 h^{-1}Mpc$ or $\approx 140 h^{-1}Mpc$ with $\delta\rho_0/\rho_0 \approx -0.11$ or $\delta\rho_0/\rho_0 \approx -0.20$ respectively, remains a viable possibility and can not be excluded by the data.

If this scenario is realized in Nature and we truly live in an underdense region, then the measured H_0 value at local scales would be larger than the true global value of H_0 . This could lead to a lower value of \mathcal{M} [Eq. (1.5)] at local scales, explaining the results of the previous section. However, a slightly off-center observer in this underdense

region would experience a preferred cosmological direction and an overall anisotropy. Therefore, in what follows we search for possible anisotropies regarding \mathcal{M} using two different methods that are widely used in the literature. These are the Hemisphere Comparison (HC) [68, 75, 88–90] and the Dipole Fitting (DF) [68, 71, 90, 91] method.

III.1. Hemisphere Comparison (HC) Method

The HC method was first proposed in Ref. [88] and implemented in the context of the Union2 dataset [92] in Ref. [75]. The basic steps of this method are the following:

- Consider a random direction of the following form

$$\hat{r}_{randm} = (\cos\phi \sqrt{1 - \cos^2\theta}, \sin\phi \sqrt{1 - \cos^2\theta}, \cos\theta) \quad (3.1)$$

where $\phi \in [0, 2\pi]$ and $\cos\theta \in [-1, 1]$. These variables are randomly selected in these intervals with a uniform probability distribution.

- Define two different hemispheres dubbed “up hemisphere” and “down hemisphere” and append the data of the dataset into the hemispheres. The appended data of the “up hemisphere” correspond to the subset where the product $\hat{r}_{randm} \cdot \hat{r}_{data}$ is positive, while the appended data of the “down hemisphere” correspond to the subset where the product $\hat{r}_{randm} \cdot \hat{r}_{data}$ is negative. The unit vector \hat{r}_{data} describes the direction of each SNIa in galactic coordinates.
- Find the best fit value of \mathcal{M} in the up (\mathcal{M}_{up}) and down hemispheres (\mathcal{M}_{down}) applying the maximum likelihood method for $\Omega_{0m} = 0.285$, *i.e.* setting Ω_{0m} to the best fit value indicated by the full dataset. Using the obtained best fit values of \mathcal{M} , define the anisotropy level (AL) as [75]

$$\Delta\mathcal{M}/\bar{\mathcal{M}} \equiv 2 \frac{\mathcal{M}_{up} - \mathcal{M}_{down}}{\mathcal{M}_{up} + \mathcal{M}_{down}} \quad (3.2)$$

as well as the corresponding 1σ error [75]

$$\sigma_{\Delta\mathcal{M}/\bar{\mathcal{M}}} = \frac{\sqrt{\sigma_{\mathcal{M}_{up}}^2 + \sigma_{\mathcal{M}_{down}}^2}}{\mathcal{M}_{up} + \mathcal{M}_{down}} \quad (3.3)$$

- Repeat this procedure for N random directions \hat{r}_{randm} and find the maximum AL and the related direction. The number of random directions needs to be well above the number of datapoints in each hemisphere [75], so for the Pantheon data we set $N = 3000$.

Implementing the HC method in the Pantheon dataset as described above, we construct the AL color map of \mathcal{M}




FIG. 6: The AL color map constructed with the HC method producing 3000 random directions. The red dots correspond to the pair of coordinates where the ratio $\Delta\mathcal{M}/\bar{\mathcal{M}}$ is maximum while the purple dots correspond to the pair of coordinates where $\Delta\mathcal{M}/\bar{\mathcal{M}}$ is minimum. The black ellipses denote the 1σ error region.




FIG. 7: *Left Panel:* The 30 axes of extrema of AL constructed from the isotropic simulated Pantheon datasets axes using 3000 random hemisphere directions in each dataset. Notice that only two of the thirty maxima AL directions are in the lower left quadrisphere (southern hemisphere in the longitude range of $[0^\circ, 180^\circ]$), inducing an artificial region of preferred directions in the observed anisotropy in the lower left/upper right quadrisphere. The green dot corresponds to the maximum anisotropy of the real data, while the blue (red) dots describe the simulated datasets which have smaller (larger) magnitudes of $\Delta\mathcal{M}/\bar{\mathcal{M}}$ than the real data. *Right Panel:* The distribution of the full Pantheon data in galactic coordinates. Notice that the data are not uniformly distributed with strong preference of datapoint locations in the southern hemisphere in the longitude range $[0^\circ, 180^\circ]$ (lower left hemisphere).

as it is demonstrated in Fig. 6. The magnitude of the maximum AL that is detected for the Pantheon data is

$$(\Delta\mathcal{M}/\bar{\mathcal{M}})_{max} = 0.0018 \pm 0.0002 \quad (3.4)$$

and the direction of the maximum anisotropy is in $(l, b) = (286.93^\circ \pm 18.52^\circ, 27.02^\circ \pm 6.50^\circ)$

In order to check the consistency of the Pantheon SNIa data with statistical isotropy we compare the above extrema of AL of the real data with the corresponding extrema of AL derived in the context of simulated Pantheon data. The simulated Pantheon data are constructed under the assumption of statistical isotropy with

a Λ CDM background by keeping fixed the direction of each datapoint in the sky while randomly selecting the Pantheon apparent magnitudes from a gaussian distribution with the best fit Λ CDM mean and standard deviation equal to the corresponding Pantheon datapoint 1σ error. We thus construct 30 isotropic simulated “Pantheon” datasets and for each dataset we use 3000 random directions to split it in two hemispheres and identify the corresponding extrema of AL using $\Delta\mathcal{M}/\bar{\mathcal{M}}$. These 30 axes of extrema of AL are shown in Fig. 7 using galactic coordinates and showing two opposite points for each maximum AL direction (left panel) along with the cor-




FIG. 8: *Left Panel:* The 100 axes of extrema of AL using 1500 random directions for each isotropically distributed Pantheon subsample. **The preferred direction disappears completely for the more isotropic distributed subset.** The green dot corresponds to the maximum anisotropy of the real data, while the blue (red) dots describe the simulated datasets which have smaller (larger) magnitudes of $\Delta\mathcal{M}/\bar{\mathcal{M}}$ than the real data. *Right Panel:* The distribution of the reduced isotropic subset in galactic coordinates.

responding real Pantheon data sky directions. The maximum AL of $\Delta\mathcal{M}/\bar{\mathcal{M}}$ magnitude of 16 (red dots in the left panel of Fig. 7) out of the 30 simulated datasets was larger than the corresponding magnitude of the real Pantheon data. **This indicates that there is no statistically significant $\Delta\mathcal{M}/\bar{\mathcal{M}}$ AL in the Pantheon data.**

Notice however, that the 30 extrema AL directions of the isotropic Pantheon simulated data are not distributed uniformly. This is due to the fact that the Pantheon SNIa are not isotropically distributed in the sky. As shown in Fig. 7 (right panel) the southern (lower) right quadrisphere is almost empty of SNIa datapoints while most of the Pantheon SNIa directions are concentrated in the southern left quadrisphere. This strongly anisotropic distribution of datapoints forces most of the extrema AL directions to concentrate in the southern left - northern right quadrisphere.

It is straightforward to reduce this effect of anisotropic distribution of Pantheon SNIa by selecting a more isotropic subset of the full dataset which will be less biased in the selection of the maximum AL direction. Thus, **we randomly select a subsample of the Pantheon dataset consisting of 375 SNIa distributed more isotropically in the four quadrispheres (100 in the first three and 75 in the down right quadrisphere) and generate a new reduced dataset (right panel of Fig. 8).** Using this reduced dataset, which is significantly more homogeneous than the full dataset, we produce 100 simulated Pantheon isotropic subsamples using 1500 random directions to split it in two hemispheres and identify the corresponding maximum $\Delta\mathcal{M}/\bar{\mathcal{M}}$ AL magnitudes². This is illustrated in the left panel of Fig. 8 where we show two

opposite points for each maximum AL direction. Clearly, the preferred range of directions disappears completely for the more isotropic subset of the full dataset. However, even in this case where the data are more uniformly distributed, no signal of anisotropy is found, since 33 (red dots in the left panel of Fig. 8) out of the 100 simulated datasets have larger maximum AL magnitudes of $\Delta\mathcal{M}/\bar{\mathcal{M}}$ than the corresponding magnitude of the real Pantheon data.

The lack of anisotropy signal persists also if we divide the Pantheon data in four redshift bins. Using the same method as described above we construct for each bin 30 isotropic simulated ‘‘Pantheon’’ datasets and for each dataset we use 1000 random directions to split the sky in two hemispheres and identify the corresponding maximum $\Delta\mathcal{M}/\bar{\mathcal{M}}$ magnitudes. Then we compare the maximum magnitudes $\Delta\mathcal{M}/\bar{\mathcal{M}}$ of the simulated ‘‘Pantheon’’ datasets with the corresponding maximum magnitude $\Delta\mathcal{M}/\bar{\mathcal{M}}$ of the real data for each bin. The results for each bin are presented in the following Table I.

TABLE I: The results of the HC method for each bin

Bin	Redshift Range	Number of Simulated Datasets with $ \frac{\Delta\mathcal{M}}{\bar{\mathcal{M}}} _{sim} > \frac{\Delta\mathcal{M}}{\bar{\mathcal{M}}} _{real}$
1st	$0.01 < z < 0.13$	21/30
2nd	$0.13 < z < 0.25$	8/30
3rd	$0.25 < z < 0.42$	14/30
4th	$0.42 < z < 2.26$	3/30

Interestingly, the strongest evidence for anisotropy is not found in the lowest z bin but in the highest z bin ($0.42 < z < 2.3$), where only three out of the thirty simulated datasets have larger $\Delta\mathcal{M}/\bar{\mathcal{M}}$ magnitudes than the corresponding magnitude of the real data. However, this mild effect is not statistically significant, since it remains below the 2σ level.

² The number of the random directions considered for the identification of the direction of the maximum AL is smaller in this case, since the new dataset is significantly smaller than the original.

Even though we have found no evidence for anisotropy of $\Delta\mathcal{M}/\mathcal{M}$ in the Pantheon data, the local underdensity scenario as an explanation for the reduced by about 4% low z value of \mathcal{M} pointed out in the previous section, remains viable especially if we are located close to the center of the underdensity. Using Eq. (2.6), we can also constrain the density contrast $\delta\rho_0/\rho_0$ as well as the dimensionless mater density contrast $\delta\Omega_0/\Omega_0$, through the following coupled system of equations applicable in the context of a LTB model with a cosmological constant assuming a top hat density profile for the void (see Appendix of Ref. [58])

$$\frac{\delta H_0}{H_0} = \frac{\delta\rho_0}{\rho_0} [-0.171 - 0.322(\Omega_{0m} - 0.3) + 0.249(\Omega_{0m} - 0.3)^2] + \left(\frac{\delta\rho_0}{\rho_0}\right)^2 [0.031 + 0.063(\Omega_{0m} - 0.3)] - 0.022 \left(\frac{\delta\rho_0}{\rho_0}\right)^3 \quad (3.5)$$

$$\frac{\delta\Omega_0}{\Omega_0} = \frac{\delta\rho_0}{\rho_0} [1.342 + 0.643(\Omega_{0m} - 0.3) - 0.499(\Omega_{0m} - 0.3)^2] + \left(\frac{\delta\rho_0}{\rho_0}\right)^2 [0.367 + 0.847(\Omega_{0m} - 0.3)] + 0.056 \left(\frac{\delta\rho_0}{\rho_0}\right)^3 \quad (3.6)$$

where in this case we set $\Omega_{0m} = 0.3153$, *i.e.* the CMB value indicated by the Planck mission [6]. Substituting $\delta H_0/H_0 \approx 0.02$ [as indicated from Eq. (2.6)] in Eq. (3.5), we calculate $\delta\rho_0/\rho_0 = -0.10 \pm 0.04$ and using this value to Eq. (3.6) we derive $\delta\Omega_0/\Omega_0 = -0.12 \pm 0.02$, in agreement with previous studies [58].

III.2. Dipole Fitting (DF) Method

In most physical mechanisms the predicted cosmological anisotropy can be described by a dipole proportional to $\cos\theta$. In this case the Dipole Fitting (DF) method [68, 71, 90, 91] is expected to be more sensitive for the detection of the cosmic anisotropy. In this context we define the deviation of the apparent magnitude from its best fit Λ CDM values $\bar{m}(z)$ as

$$\left(\frac{\Delta m(z)}{\bar{m}(z)}\right)_{obs} \equiv \frac{\bar{m}(z) - m(z)}{\bar{m}(z)} \quad (3.7)$$

The basic steps of the DF method are the following [91]

- Convert the coordinates of the SnIa to galactic coordinates (l, b) (they are provided in equatorial coordinates) and define the unit vector \hat{n}_i as

$$\hat{n}_i = \cos(b_i) \cos(l_i) \hat{x} + \cos(b_i) \sin(l_i) \hat{y} + \sin(b_i) \hat{z} \quad (3.8)$$

- Define the dipole axis \vec{D} in terms of the parameters c_1, c_2 and c_3 in cartesian coordinates as

$$\vec{D} = c_1 \hat{x} + c_2 \hat{y} + c_3 \hat{z} \quad (3.9)$$

and define

$$\left(\frac{\Delta m}{\bar{m}}\right)_{th} = A \cos\theta + B \quad (3.10)$$

where A and B correspond to the dipole and monopole terms of the parametrized anisotropy. The angle θ is the angle between the datapoint direction of the SnIa with the vector \vec{D} which obeys the relation

$$\hat{n}_i \vec{D} = A \cos\theta_i \quad (3.11)$$

- Using the maximum likelihood method construct χ^2 as

$$\chi^2 = \sum_{i=1}^{1048} V^i C_{ij}^{-1} V^j \quad (3.12)$$

where $V^i \equiv (\Delta m/m)_{obs} - (\Delta m/m)_{th} = [\bar{m}(z_i) - m(z_i)]/\bar{m}(z_i) - A \cos\theta_i - B$ and C_{ij} is the covariance matrix. From the minimization of χ^2 , find the best fit values as well as the 1σ errors of the c_i , the monopole term B and thus the dipole term that is derived as $A = \sqrt{\sum_{j=1}^3 c_j^2}$.

We present the results of the application of the DF method for the Pantheon dataset in the following Table II.

TABLE II: The best fit values with the 1σ error of the c_i 's, A and B parameters.

Quantity	Best Fit Value $\pm 1\sigma$ Error
c_1	$(-1.41 \pm 3.76) \times 10^{-4}$
c_2	$(-0.82 \pm 4.54) \times 10^{-4}$
c_3	$(5.28 \pm 7.14) \times 10^{-4}$
A	$(5.53 \pm 6.04) \times 10^{-4}$
B	$(-0.59 \pm 3.01) \times 10^{-4}$
l	$210.254^\circ \pm 136.564$
b	$72.852^\circ \pm 60.631^\circ$

The monopole A and dipole B terms are consistent with zero at the 1σ level. Using the best fit values of the parameters A and B , we find the anisotropy direction to be $(l, b) = (210.254^\circ \pm 136.564, 72.852^\circ \pm 60.631^\circ)$. Clearly, the errors of the (l, b) coordinates are quite large, covering almost the entire sky area (a result that is in agreement with previous studies [68–70] using Ω_{0m} instead of \mathcal{M}).

The consistency of the derived dipole and monopole terms with statistical isotropy may be investigated using isotropic simulated Pantheon datasets as we did in the context of the HC method. We construct 30 simulated Pantheon datasets as described in the previous subsection, we identify the corresponding dipole anisotropy directions and the best fit values of the parameters A and B as shown in Fig. 9.




FIG. 9: The different maximum AL directions corresponding to the 30 random simulated datasets. The green dot corresponds to dipole of the real data, while the blue (red) dots describe the dipole direction of simulated datasets which have smaller (larger) magnitudes of A than the real data. The 1σ errors of the (l, b) galactic coordinates are quite large covering almost the entire sky area.

From Fig. 9, it is clear that no preferred direction is identified since 19 (red points in Fig. 9) of the 30 isotropic simulated Pantheon datasets have larger dipole magnitudes than the real data. Therefore, we conclude that no statistically significant anisotropy is found using the DF method, in agreement with the corresponding result of the HC method.

III.3. Comparison of the Two Methods

From the implementation of the two methods the following useful conclusions, can be extracted

- The HC method is more general, since it can detect any kind of anisotropy. On the contrary, the DF method is sensitive only to an anisotropy of the form of Eq. (3.10), *i.e.* an anisotropy that has a dipole form.
- The 1σ errors of the anisotropy direction coordinates obtained in the context of the DF method are quite large and cover the entire sky area. Thus, a dipole anisotropy seems to be significantly disfavoured by the Pantheon dataset indicating that no dipole signal exists in the data. On the contrary, the HC method gives significantly smaller 1σ errors as it is tuned for the detection of a much broader range of signals. Therefore, the HC method seems to be more appropriate in order to identify a preferred direction as well as any general anisotropies hidden in the Pantheon data, unless these anisotropies are of the particular dipole form.

In conclusion, we have found no evidence of anisotropy in the Pantheon data, a result consistent with previous studies [66–70, 93]. We have shown however, that the HC method is more appropriate in detecting a general form

of anisotropy hidden in the data. We have also demonstrated that the anisotropic distribution of the Pantheon SNIa data leads to a preferred range of anisotropy directions which are detected by the HC method in the context of isotropic simulated Pantheon datasets. This lack of anisotropy does not favour (but also does not exclude) the local underdensity scenario as a possible explanation of the observed reduced value of \mathcal{M} at low z indicated in section II. We thus proceed to examine the alternative mechanism that could lead to a reduced value of \mathcal{M} at low z : the evolving μ scenario.

IV. MODIFIED THEORY OF GRAVITY SCENARIO

In order to identify the possible evolution of μ we consider Eq. (2.4) and use the 100 point moving subsample method described in Section II (Figure 2). In particular, we find the best fit value of M , fixing M_0 to the best fit value of the absolute magnitude M indicated by the full dataset with $h = 0.74$. Then, for $b = -3/2$, we use the best fit values of M and find the corresponding best fit values of μ , thus assigning any redshift dependence of M into a redshift dependence of μ . The resulting best fit values of μ for each subsample along with the 1σ errors are shown in Fig. 10.




FIG. 10: The evolution of μ along with its 1σ error vs the mean redshift z_{mean} of each 100 datapoints subsample. In the context a modified theory, we detect a 2 – 3% deviation from the GR predicted value $\mu = 1$ (dashed line) at a level up to about 2σ .

Clearly the oscillating behaviour of M at low z shown in Fig. 2 middle panel, is reflected on a corresponding oscillating behaviour for μ at low z with $\mu < 1$ at $z = 0$.

We now consider a μ parametrization which interpolates GR at early and late times and takes into consideration the solar system and nucleosynthesis constraints. This parametrization is the following [44, 45]

$$\mu(z, g_a) = 1 + g_a \left(\frac{z}{1+z} \right)^2 - g_a \left(\frac{z}{1+z} \right)^4 \quad (4.1)$$

where g_a is an extra parameter and z is the redshift. Using the modified apparent magnitude (2.4) along with the parametrization (4.1) we construct the corresponding χ^2 function and applying the maximum likelihood method, we obtain the best fit values for the parameters $\mathcal{M}, \Omega_{0m}, g_a$ and b . In this case, χ^2 depends on the same parameters as before (\mathcal{M}, Ω_{0m}) as well as the extra parameters g_a and b . During the minimization we allow b to take various values in the range $-2 < b < 2$ and interpolate the best fits of the extra parameter g_a as a function of b . This is demonstrated in Fig. 11.




FIG. 11: The extra parameter g_a as a function of b . For $b > 0$ we obtain $g_a > 0$ while for $b < 0$ we find a best fit $g_a < 0$.

For negative values of b we obtain negative best fit values for g_a indicating that $\mu < 1$, *i.e.* a growth rate that is weaker than expected in the context of Λ CDM. This result is in agreement with other studies [44–46, 48] which fit weak lensing and growth data allowing for an evolving μ and favour an evolving μ with $g_a < 0$.

Setting $b = -3/2$ as indicated by most relevant studies [63, 64] it is straightforward to construct the likelihood parameter contours in the 3D parametric space $(\mathcal{M}, \Omega_{0m}, g_a)$ where \mathcal{M} is given by Eq. (1.5). The best fit parameter values thus obtained are $\mathcal{M} = 23.793 \pm 0.009$, $\Omega_{0m} = 0.179 \pm 0.078$ and $g_a = -0.47 \pm 0.36$. In Fig. 12 we show the 2D projections of the $1\sigma - 4\sigma$ contours in the parametric space $(\mathcal{M}, \Omega_{0m}, g_a)$. The projections go through the best fit point in the 3D parameter space. Notice that the GR point corresponding to $g_a = 0$ appears to be more than 4σ away from the best fit which corresponds to weaker gravity ($g_a < 0$) in accordance with weak lensing and growth cosmological data [33–36, 44–46, 48]. However, this is a projection effect since in the context of the full 3D parameter space we have $g_a = -0.47 \pm 0.36$, a value that is approximately 1.5σ away from the GR predicted one.

V. CONCLUSION - DISCUSSION - OUTLOOK

We have performed a redshift tomographic analysis of the latest SnIa (Pantheon) data in the context of a Λ CDM model fitting simultaneously the matter density parameter Ω_{0m} and the parameter \mathcal{M} which depends on both the calibrated absolute magnitude M and the Hubble parameter H_0 . We have found a mild tension ($2 - 3\sigma$) between the best fit value of \mathcal{M} obtained from low z SnIa ($z \in [0.01, 0.2]$) and the corresponding value obtained from the full Pantheon dataset. If this tension is not a statistical fluctuation or some unknown systematic effect in the data it could be either explained as a locally higher value of H_0 corresponding to a local underdensity with $(\frac{\delta\rho_0}{\rho_0}, \frac{\delta\Omega_0}{\Omega_0}) \simeq (-0.10 \pm 0.04, -0.12 \pm 0.02)$ or as a modified gravity effect leading to a time variation of Newton’s constant.

In the context of the local underdensity scenario, a degree of anisotropy is anticipated for \mathcal{M} , unless the observer is located at the center of this underdensity. Thus we used two methods to search for statistically significant anisotropy in the Pantheon SnIa data: The Hemisphere Comparison (HC) method and the Dipole Fit (DF) method. Even though we found no statistically significant evidence for cosmological anisotropy our analysis revealed the following interesting facts:

- Using simulated Pantheon-like data constructed under the assumption of an underlying Λ CDM model, we showed that the anisotropic distribution of the SnIa datapoints in the sky generically favours the range anisotropy of directions consistent with the dataset in $b \in [-15^\circ, -45^\circ]$, $l \in [60^\circ, 150^\circ]$ (or in the opposite direction $b \in [15^\circ, 45^\circ]$, $l \in [240^\circ, 330^\circ]$). We constructed a more isotropically distributed subset of the Pantheon data that appears to be free of this limitation but less powerful in detecting overall anisotropy due to reduced number of datapoints.
- The HC method appears to be more powerful in detecting a general anisotropy signal than the DF method since the statistical uncertainties for both the magnitude and direction of anisotropy appear to be smaller in the context of this method.

The lack of evidence for statistical anisotropy in the SnIa data does not favour the local underdensity scenario as a possible explanation for the reduced value of \mathcal{M} at low z as such anisotropy would be expected in the context of a non-spherical anisotropy and/or an off-center observer.

The abnormal variation of \mathcal{M} can also be explained in the context of a modified theory of gravity with an evolving Newton’s constant at low redshifts. In this context, allowing for an evolving normalized Newton’s constant $\mu(z)$, we found a 2σ deviation of $\sim 2 - 3\%$ from the GR predicted value at low redshifts. Moreover, considering a physically motivated parametrization for the evolving Newton’s constant that interpolates GR at early and late




FIG. 12: The 2D projections of the $1\sigma - 4\sigma$ contours in the parametric space $(\mathcal{M}, \Omega_{0m}, g_a)$. The projections go through the best fit point (green point) in the 3D parameter space. Notice that the GR point corresponding to $g_a = 0$ appears to be more than 4σ away from the best fit which corresponds to weaker gravity ($g_a < 0$). However, this is a projection effect since in the context of the full 3D parameter space we have $g_a = -0.47 \pm 0.36$.

times [Eq. (4.1)], we derived the best fit value of the extra parameter g_a as $g_a = -0.47 \pm 0.36$. This value is approximately 1.5σ away from the GR predicted value ($g_a = 0$), favouring a reduced Newton’s constant compared to GR. This weak hint is consistent with the results at low z of other studies that mildly favour weakening gravity using growth [44, 45, 47, 48] and weak lensing data [33–36].

Interesting extensions of the present analysis include the following:

- Use of an extended up to date SnIa dataset with more uniform distribution in the sky to investigate and further constrain the possible evolution of the parameter \mathcal{M} with redshift and its connection with a possible cosmic anisotropy.
- Further investigate the connection between an evolving \mathcal{M} and an evolving Newton’s constant in the context of various models for the mechanism of SnIa explosion. In particular a reliable estimate of the sign and value of the power index b that connects the evolving absolute magnitude M with the effective Newton’s constant is important for imposing reliable constraints on modified gravity models from the possible evolution of SnIa absolute luminosity.
- The use of alternative standard candle probes (e.g. γ ray bursts) to search for possible similar hints of

variation of H_0 and/or G_{eff} .

- The identification of new statistical tests probing for cosmological anisotropies of SnIa data and the comparison of their efficiency with the standard methods used in the present analysis (HC and DF).
- The consideration of alternative background expansion cosmologies. It may be possible to absorb the variation of \mathcal{M} at low z in the context of a varying dark energy equation of state parameter w at low z . In this context, a varying \mathcal{M} at low z may be a hint for a variation of w . Such variation may also play a role in resolving the H_0 problem [94–96].

Numerical Analysis Files: The numerical files for the reproduction of the figures can be found in [97].

ACKNOWLEDGEMENTS

This research is co-financed by Greece and the European Union (European Social Fund- ESF) through the Operational Programme “Human Resources Development, Education and Lifelong Learning” in the context of the project “Strengthening Human Resources Research Potential via Doctorate Research – 2nd Cycle” (MIS-5000432), implemented by the State Scholarships Foundation (IKY).

[1] Adam G. Riess *et al.* (Supernova Search Team), “Observational evidence from supernovae for an accelerating universe and a cosmological constant,” *Astron. J.* **116**, 1009–1038 (1998), arXiv:astro-ph/9805201 [astro-ph].

[2] S. Perlmutter *et al.* (Supernova Cosmology Project), “Measurements of Ω and Λ from 42 high redshift supernovae,” *Astrophys. J.* **517**, 565–586 (1999), arXiv:astro-ph/9812133 [astro-ph].

- [3] Sean M. Carroll, “The Cosmological constant,” *Living Rev. Rel.* **4**, 1 (2001), arXiv:astro-ph/0004075 [astro-ph].
- [4] G. Hinshaw *et al.* (WMAP), “Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Parameter Results,” *Astrophys. J. Suppl.* **208**, 19 (2013), arXiv:1212.5226 [astro-ph.CO].
- [5] P. A. R. Ade *et al.* (Planck), “Planck 2015 results. XIII. Cosmological parameters,” *Astron. Astrophys.* **594**, A13 (2016), arXiv:1502.01589 [astro-ph.CO].
- [6] N. Aghanim *et al.* (Planck), “Planck 2018 results. VI. Cosmological parameters,” (2018), arXiv:1807.06209 [astro-ph.CO].
- [7] M. Betoule *et al.* (SDSS), “Improved cosmological constraints from a joint analysis of the SDSS-II and SNLS supernova samples,” *Astron. Astrophys.* **568**, A22 (2014), arXiv:1401.4064 [astro-ph.CO].
- [8] D. M. Scolnic *et al.*, “The Complete Light-curve Sample of Spectroscopically Confirmed SNe Ia from Pan-STARRS1 and Cosmological Constraints from the Combined Pantheon Sample,” *Astrophys. J.* **859**, 101 (2018), arXiv:1710.00845 [astro-ph.CO].
- [9] Daniel Stern, Raul Jimenez, Licia Verde, Marc Kamionkowski, and S. Adam Stanford, “Cosmic Chronometers: Constraining the Equation of State of Dark Energy. I: H(z) Measurements,” *JCAP* **1002**, 008 (2010), arXiv:0907.3149 [astro-ph.CO].
- [10] M. Moresco *et al.*, “Improved constraints on the expansion rate of the Universe up to z 1.1 from the spectroscopic evolution of cosmic chronometers,” *JCAP* **1208**, 006 (2012), arXiv:1201.3609 [astro-ph.CO].
- [11] Adrià Gómez-Valent and Luca Amendola, “ H_0 from cosmic chronometers and Type Ia supernovae, with Gaussian Processes and the novel Weighted Polynomial Regression method,” *JCAP* **1804**, 051 (2018), arXiv:1802.01505 [astro-ph.CO].
- [12] Shadab Alam *et al.* (BOSS), “The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological analysis of the DR12 galaxy sample,” *Mon. Not. Roy. Astron. Soc.* **470**, 2617–2652 (2017), arXiv:1607.03155 [astro-ph.CO].
- [13] Éric Aubourg *et al.*, “Cosmological implications of baryon acoustic oscillation measurements,” *Phys. Rev.* **D92**, 123516 (2015), arXiv:1411.1074 [astro-ph.CO].
- [14] Spyros Basilakos and Savvas Nesseris, “Testing Einstein’s gravity and dark energy with growth of matter perturbations: Indications for new physics?” *Phys. Rev.* **D94**, 123525 (2016), arXiv:1610.00160 [astro-ph.CO].
- [15] Alba Quelle and Antonio L. Maroto, “On the tension between growth rate and CMB data,” (2019), arXiv:1908.00900 [astro-ph.CO].
- [16] E. Baxter *et al.*, “Joint measurement of lensing–galaxy correlations using SPT and DES SV data,” *Mon. Not. Roy. Astron. Soc.* **461**, 4099–4114 (2016), arXiv:1602.07384 [astro-ph.CO].
- [17] George Efstathiou and Pablo Lemos, “Statistical inconsistencies in the KiDS-450 data set,” *Mon. Not. Roy. Astron. Soc.* **476**, 151–157 (2018), arXiv:1707.00483 [astro-ph.CO].
- [18] David Rapetti, Steven W. Allen, Adam Mantz, and Harald Ebeling, “Constraints on modified gravity from the observed X-ray luminosity function of galaxy clusters,” *Mon. Not. Roy. Astron. Soc.* **400**, 699 (2009), arXiv:0812.2259 [astro-ph].
- [19] Eduardo Rozo *et al.* (DSDD), “Cosmological Constraints from the SDSS maxBCG Cluster Catalog,” *Astrophys. J.* **708**, 645–660 (2010), arXiv:0902.3702 [astro-ph.CO].
- [20] P. A. R. Ade *et al.* (Planck), “Planck 2015 results. XXIV. Cosmology from Sunyaev-Zeldovich cluster counts,” *Astron. Astrophys.* **594**, A24 (2016), arXiv:1502.01597 [astro-ph.CO].
- [21] Steven Weinberg, “The Cosmological Constant Problem,” *Rev. Mod. Phys.* **61**, 1–23 (1989), [569(1988)].
- [22] Varun Sahni, “The Cosmological constant problem and quintessence,” *The early universe and cosmological observations: A critical review. Proceedings, Conference, Cape Town, South Africa, July 23-25, 2001, Class. Quant. Grav.* **19**, 3435–3448 (2002), arXiv:astro-ph/0202076 [astro-ph].
- [23] Steinhardt P.J., *Critical Problems in Physics* (Princeton University Press, 1997).
- [24] H. E. S. Velten, R. F. vom Marttens, and W. Zimdahl, “Aspects of the cosmological “coincidence problem”,” *Eur. Phys. J.* **C74**, 3160 (2014), arXiv:1410.2509 [astro-ph.CO].
- [25] Guido Risaliti and Elisabeta Lusso, “Cosmological constraints from the Hubble diagram of quasars at high redshifts,” *Nat. Astron.* **3**, 272–277 (2019), arXiv:1811.02590 [astro-ph.CO].
- [26] Rubén Arjona and Savvas Nesseris, “What can Machine Learning tell us about the background expansion of the Universe?” (2019), arXiv:1910.01529 [astro-ph.CO].
- [27] Eleonora Di Valentino, Alessandro Melchiorri, and Joseph Silk, “Planck evidence for a closed Universe and a possible crisis for cosmology,” *Nat. Astron.* (2019), 10.1038/s41550-019-0906-9, arXiv:1911.02087 [astro-ph.CO].
- [28] Rubén Arjona and Savvas Nesseris, “Hints of dark energy anisotropic stress using Machine Learning,” (2020), arXiv:2001.11420 [astro-ph.CO].
- [29] Adam G. Riess, Stefano Casertano, Wenlong Yuan, Lucas M. Macri, and Dan Scolnic, “Large Magellanic Cloud Cepheid Standards Provide a 1% Foundation for the Determination of the Hubble Constant and Stronger Evidence for Physics beyond Λ CDM,” *Astrophys. J.* **876**, 85 (2019), arXiv:1903.07603 [astro-ph.CO].
- [30] Kenneth C. Wong *et al.*, “H0LiCOW XIII. A 2.4% measurement of H_0 from lensed quasars: 5.3 σ tension between early and late-Universe probes,” (2019), arXiv:1907.04869 [astro-ph.CO].
- [31] Adam G. Riess, “The Expansion of the Universe is Faster than Expected,” *Nature Rev. Phys.* **2**, 10–12 (2019), arXiv:2001.03624 [astro-ph.CO].
- [32] Wendy L. Freedman *et al.*, “The Carnegie-Chicago Hubble Program. VIII. An Independent Determination of the Hubble Constant Based on the Tip of the Red Giant Branch,” (2019), 10.3847/1538-4357/ab2f73, arXiv:1907.05922 [astro-ph.CO].
- [33] H. Hildebrandt *et al.*, “KiDS-450: Cosmological parameter constraints from tomographic weak gravitational lensing,” *Mon. Not. Roy. Astron. Soc.* **465**, 1454 (2017), arXiv:1606.05338 [astro-ph.CO].
- [34] F. Köhlinger *et al.*, “KiDS-450: The tomographic weak lensing power spectrum and constraints on cosmological parameters,” *Mon. Not. Roy. Astron. Soc.* **471**, 4412–4435 (2017), arXiv:1706.02892 [astro-ph.CO].
- [35] Shahab Joudaki *et al.*, “KiDS-450 + 2dFLenS: Cosmological parameter constraints from weak gravitational lens-

- ing tomography and overlapping redshift-space galaxy clustering,” *Mon. Not. Roy. Astron. Soc.* **474**, 4894–4924 (2018), arXiv:1707.06627 [astro-ph.CO].
- [36] T. M. C. Abbott *et al.* (DES), “Dark Energy Survey year 1 results: Cosmological constraints from galaxy clustering and weak lensing,” *Phys. Rev.* **D98**, 043526 (2018), arXiv:1708.01530 [astro-ph.CO].
- [37] T. M. C. Abbott *et al.* (DES), “Dark Energy Survey Year 1 Results: Constraints on Extended Cosmological Models from Galaxy Clustering and Weak Lensing,” *Phys. Rev.* **D99**, 123505 (2019), arXiv:1810.02499 [astro-ph.CO].
- [38] Edward Macaulay, Ingunn Kathrine Wehus, and Hans Kristian Eriksen, “Lower Growth Rate from Recent Redshift Space Distortion Measurements than Expected from Planck,” *Phys. Rev. Lett.* **111**, 161301 (2013), arXiv:1303.6583 [astro-ph.CO].
- [39] Andrew Johnson, Chris Blake, Jason Dossett, Jun Koda, David Parkinson, and Shahab Joudaki, “Searching for Modified Gravity: Scale and Redshift Dependent Constraints from Galaxy Peculiar Velocities,” *Mon. Not. Roy. Astron. Soc.* **458**, 2725–2744 (2016), arXiv:1504.06885 [astro-ph.CO].
- [40] Shinji Tsujikawa, “Possibility of realizing weak gravity in redshift space distortion measurements,” *Phys. Rev.* **D92**, 044029 (2015), arXiv:1505.02459 [astro-ph.CO].
- [41] B. Wang, E. Abdalla, F. Atrio-Barandela, and D. Pavon, “Dark Matter and Dark Energy Interactions: Theoretical Challenges, Cosmological Implications and Observational Signatures,” *Rept. Prog. Phys.* **79**, 096901 (2016), arXiv:1603.08299 [astro-ph.CO].
- [42] Joan Solà, “Cosmological constant vis-a-vis dynamical vacuum: bold challenging the Λ CDM,” *Int. J. Mod. Phys.* **A31**, 1630035 (2016), arXiv:1612.02449 [astro-ph.CO].
- [43] Spyros Basilakos and Savvas Nesseris, “Conjoined constraints on modified gravity from the expansion history and cosmic growth,” *Phys. Rev.* **D96**, 063517 (2017), arXiv:1705.08797 [astro-ph.CO].
- [44] Savvas Nesseris, George Pantazis, and Leandros Perivolaropoulos, “Tension and constraints on modified gravity parametrizations of $G_{\text{eff}}(z)$ from growth rate and Planck data,” *Phys. Rev.* **D96**, 023542 (2017), arXiv:1703.10538 [astro-ph.CO].
- [45] Lavrentios Kazantzidis and Leandros Perivolaropoulos, “Evolution of the $f\sigma_8$ tension with the Planck15/ Λ CDM determination and implications for modified gravity theories,” *Phys. Rev.* **D97**, 103503 (2018), arXiv:1803.01337 [astro-ph.CO].
- [46] Leandros Perivolaropoulos and Lavrentios Kazantzidis, “Hints of modified gravity in cosmos and in the lab?” *Int. J. Mod. Phys.* **D28**, 1942001 (2019), arXiv:1904.09462 [gr-qc].
- [47] Lavrentios Kazantzidis and Leandros Perivolaropoulos, “Is gravity getting weaker at low z ? Observational evidence and theoretical implications,” (2019), arXiv:1907.03176 [astro-ph.CO].
- [48] F. Skara and L. Perivolaropoulos, “Tension of the E_G statistic and redshift space distortion data with the Planck - Λ CDM model and implications for weakening gravity,” *Phys. Rev. D* **101**, 063521 (2020), arXiv:1911.10609 [astro-ph.CO].
- [49] G. E. Addison, D. J. Watts, C. L. Bennett, M. Halpern, G. Hinshaw, and J. L. Weiland, “Elucidating Λ CDM: Impact of Baryon Acoustic Oscillation Measurements on the Hubble Constant Discrepancy,” *Astrophys. J.* **853**, 119 (2018), arXiv:1707.06547 [astro-ph.CO].
- [50] Andrei Cuceu, James Farr, Pablo Lemos, and Andreu Font-Ribera, “Baryon Acoustic Oscillations and the Hubble Constant: Past, Present and Future,” *JCAP* **1910**, 044 (2019), arXiv:1906.11628 [astro-ph.CO].
- [51] G. E. Addison, Y. Huang, D. J. Watts, C. L. Bennett, M. Halpern, G. Hinshaw, and J. L. Weiland, “Quantifying discordance in the 2015 Planck CMB spectrum,” *Astrophys. J.* **818**, 132 (2016), arXiv:1511.00055 [astro-ph.CO].
- [52] N. Aghanim *et al.* (Planck), “Planck 2018 results. V. CMB power spectra and likelihoods,” (2019), arXiv:1907.12875 [astro-ph.CO].
- [53] Adam G. Riess *et al.*, “A Redetermination of the Hubble Constant with the Hubble Space Telescope from a Differential Distance Ladder,” *Astrophys. J.* **699**, 539–563 (2009), arXiv:0905.0695 [astro-ph.CO].
- [54] Matt Visser, “Jerk and the cosmological equation of state,” *Class. Quant. Grav.* **21**, 2603–2616 (2004), arXiv:gr-qc/0309109 [gr-qc].
- [55] A. Conley *et al.* (SNLS), “Supernova Constraints and Systematic Uncertainties from the First 3 Years of the Supernova Legacy Survey,” *Astrophys. J. Suppl.* **192**, 1 (2011), arXiv:1104.1443 [astro-ph.CO].
- [56] Tom Shanks, Lucy Hogarth, and Nigel Metcalfe, “Gaia Cepheid parallaxes and ‘Local Hole’ relieve H_0 tension,” *Mon. Not. Roy. Astron. Soc.* **484**, L64–L68 (2019), arXiv:1810.02595 [astro-ph.CO].
- [57] T. Shanks, L. M. Hogarth, N. Metcalfe, and J. Whitbourn, “Local Hole revisited: evidence for bulk motions and self-consistent outflow,” *Mon. Not. Roy. Astron. Soc.* **490**, 4715–4720 (2019), arXiv:1909.01878 [astro-ph.CO].
- [58] Vladimir V. Luković, Balakrishna S. Haridasu, and Nicola Vittorio, “Exploring the evidence for a large local void with supernovae Ia data,” *Mon. Not. Roy. Astron. Soc.* **491**, 2075–2087 (2020), arXiv:1907.11219 [astro-ph.CO].
- [59] Georges Lemaître, “L’Univers en expansion,” *Annales de la Sociéacutet&ecute; Scientifique de Bruxelles* **53**, 51 (1933).
- [60] Richard C. Tolman, “Effect of inhomogeneity on cosmological models,” *Proc. Nat. Acad. Sci.* **20**, 169–176 (1934), [Gen. Rel. Grav.29,935(1997)].
- [61] H. Bondi, “Spherically symmetrical models in general relativity,” *Mon. Not. Roy. Astron. Soc.* **107**, 410–425 (1947).
- [62] Hans Boehringer, Gayoung Chon, and Chris A. Collins, “Observational evidence for a local underdensity in the Universe and its effect on the measurement of the Hubble Constant,” *Astron. Astrophys.* **633**, A19 (2020), arXiv:1907.12402 [astro-ph.CO].
- [63] E. Garcia-Berro, E. Gaztanaga, J. Isern, O. Benvenuto, and L. Althaus, “On the evolution of cosmological type ia supernovae and the gravitational constant,” (1999), arXiv:astro-ph/9907440 [astro-ph].
- [64] E. Gaztanaga, E. Garcia-Berro, J. Isern, E. Bravo, and I. Dominguez, “Bounds on the possible evolution of the gravitational constant from cosmological type Ia supernovae,” *Phys. Rev.* **D65**, 023506 (2002), arXiv:astro-ph/0109299 [astro-ph].
- [65] Bill S. Wright and Baojiu Li, “Type Ia supernovae, standardizable candles, and gravity,” *Phys. Rev.* **D97**, 083505 (2018), arXiv:1710.07018 [astro-ph.CO].

- [66] Z. Q. Sun and F. Y. Wang, “Testing the anisotropy of cosmic acceleration from Pantheon supernovae sample,” *Mon. Not. Roy. Astron. Soc.* **478**, 5153–5158 (2018), arXiv:1805.09195 [astro-ph.CO].
- [67] Uendert Andrade, Carlos A. P. Bengaly, Beethoven Santos, and Jailson S. Alcaniz, “A Model-independent Test of Cosmic Isotropy with Low- z Pantheon Supernovae,” *Astrophys. J.* **865**, 119 (2018), arXiv:1806.06990 [astro-ph.CO].
- [68] Hua-Kai Deng and Hao Wei, “Null signal for the cosmic anisotropy in the Pantheon supernovae data,” *Eur. Phys. J.* **C78**, 755 (2018), arXiv:1806.02773 [astro-ph.CO].
- [69] Dong Zhao, Yong Zhou, and Zhe Chang, “Anisotropy of the Universe via the Pantheon supernovae sample revisited,” *Mon. Not. Roy. Astron. Soc.* **486**, 5679–5689 (2019), arXiv:1903.12401 [astro-ph.CO].
- [70] Zhe Chang, Dong Zhao, and Yong Zhou, “Constraining the anisotropy of the Universe via Pantheon supernovae sample,” (2019), arXiv:1910.06883 [astro-ph.CO].
- [71] Hai-Nan Lin, Sai Wang, Zhe Chang, and Xin Li, “Testing the isotropy of the Universe by using the JLA compilation of type-Ia supernovae,” *Mon. Not. Roy. Astron. Soc.* **456**, 1881–1885 (2016), arXiv:1504.03428 [astro-ph.CO].
- [72] Hua-Kai Deng and Hao Wei, “Testing the Cosmic Anisotropy with Supernovae Data: Hemisphere Comparison and Dipole Fitting,” *Phys. Rev.* **D97**, 123515 (2018), arXiv:1804.03087 [astro-ph.CO].
- [73] Z. Q. Sun and F. Y. Wang, “Probing the isotropy of cosmic acceleration using different supernova samples,” *Eur. Phys. J.* **C79**, 783 (2019), arXiv:1804.05191 [astro-ph.CO].
- [74] Rubén Arjona, Wilmar Cardona, and Savvas Nesseris, “Unraveling the effective fluid approach for $f(R)$ models in the subhorizon approximation,” *Phys. Rev.* **D99**, 043516 (2019), arXiv:1811.02469 [astro-ph.CO].
- [75] I. Antoniou and L. Perivolaropoulos, “Searching for a Cosmological Preferred Axis: Union2 Data Analysis and Comparison with Other Probes,” *JCAP* **1012**, 012 (2010), arXiv:1007.4347 [astro-ph.CO].
- [76] Eoin Ó. Colgáin, “A hint of matter underdensity at low z ?” *JCAP* **1909**, 006 (2019), arXiv:1903.11743 [astro-ph.CO].
- [77] Havard Alnes, Morad Amarzguoui, and Oyvind Gron, “An inhomogeneous alternative to dark energy?” *Phys. Rev.* **D73**, 083519 (2006), arXiv:astro-ph/0512006 [astro-ph].
- [78] Juan Garcia-Bellido and Troels Haugboelle, “Confronting Lemaitre-Tolman-Bondi models with Observational Cosmology,” *JCAP* **0804**, 003 (2008), arXiv:0802.1523 [astro-ph].
- [79] Kari Enqvist and Teppo Mattsson, “The effect of inhomogeneous expansion on the supernova observations,” *JCAP* **0702**, 019 (2007), arXiv:astro-ph/0609120 [astro-ph].
- [80] Sean February, Julien Larena, Mathew Smith, and Chris Clarkson, “Rendering Dark Energy Void,” *Mon. Not. Roy. Astron. Soc.* **405**, 2231 (2010), arXiv:0909.1479 [astro-ph.CO].
- [81] Tirthabir Biswas, Alessio Notari, and Wessel Valkenburg, “Testing the Void against Cosmological data: fitting CMB, BAO, SN and H_0 ,” *JCAP* **1011**, 030 (2010), arXiv:1007.3065 [astro-ph.CO].
- [82] Hao-Yi Wu and Dragan Huterer, “Sample variance in the local measurements of the Hubble constant,” *Mon. Not. Roy. Astron. Soc.* **471**, 4946–4955 (2017), arXiv:1706.09723 [astro-ph.CO].
- [83] William J. Frith, G. S. Busswell, R. Fong, N. Metcalfe, and T. Shanks, “The local hole in the galaxy distribution: Evidence from 2MASS,” *Mon. Not. Roy. Astron. Soc.* **345**, 1049 (2003), arXiv:astro-ph/0302331 [astro-ph].
- [84] William J. Frith, N. Metcalfe, and T. Shanks, “New h-band galaxy number counts: a large local hole in the galaxy distribution?” *Mon. Not. Roy. Astron. Soc.* **371**, 1601–1609 (2006), arXiv:astro-ph/0509875 [astro-ph].
- [85] Ryan C. Keenan, Amy J. Barger, and Lennox L. Cowie, “Evidence for a 300 Megaparsec Scale Under-density in the Local Galaxy Distribution,” *Astrophys. J.* **775**, 62 (2013), arXiv:1304.2884 [astro-ph.CO].
- [86] J. R. Whitbourn and T. Shanks, “The Local Hole revealed by galaxy counts and redshifts,” *Mon. Not. Roy. Astron. Soc.* **437**, 2146–2162 (2014), arXiv:1307.4405 [astro-ph.CO].
- [87] J. R. Whitbourn and T. Shanks (Extragalactic Astronomy Group, Durham University), “The galaxy luminosity function and the Local Hole,” *Mon. Not. Roy. Astron. Soc.* **459**, 496–507 (2016), arXiv:1603.02322 [astro-ph.CO].
- [88] Dominik J. Schwarz and Bastian Weinhorst, “(An)isotropy of the Hubble diagram: Comparing hemispheres,” *Astron. Astrophys.* **474**, 717–729 (2007), arXiv:0706.0165 [astro-ph].
- [89] Rong-Gen Cai and Zhong-Liang Tuo, “Direction Dependence of the Deceleration Parameter,” *JCAP* **1202**, 004 (2012), arXiv:1109.0941 [astro-ph.CO].
- [90] Zhe Chang and Hai-Nan Lin, “Comparison between hemisphere comparison method and dipole-fitting method in tracing the anisotropic expansion of the Universe use the Union2 dataset,” *Mon. Not. Roy. Astron. Soc.* **446**, 2952–2958 (2015), arXiv:1411.1466 [astro-ph.CO].
- [91] Antonio Mariano and Leandros Perivolaropoulos, “Is there correlation between Fine Structure and Dark Energy Cosmic Dipoles?” *Phys. Rev.* **D86**, 083517 (2012), arXiv:1206.4055 [astro-ph.CO].
- [92] R. Amanullah *et al.*, “Spectra and Light Curves of Six Type Ia Supernovae at 0.511 $\leq z \leq$ 1.12 and the Union2 Compilation,” *Astrophys. J.* **716**, 712–738 (2010), arXiv:1004.1711 [astro-ph.CO].
- [93] Daniela Saadeh, Stephen M. Feeney, Andrew Pontzen, Hiranya V. Peiris, and Jason D. McEwen, “How isotropic is the Universe?” *Phys. Rev. Lett.* **117**, 131302 (2016), arXiv:1605.07178 [astro-ph.CO].
- [94] Weiqiang Yang, Supriya Pan, Eleonora Di Valentino, Emmanuel N. Saridakis, and Subenoy Chakraborty, “Observational constraints on one-parameter dynamical dark-energy parametrizations and the H_0 tension,” *Phys. Rev.* **D99**, 043543 (2019), arXiv:1810.05141 [astro-ph.CO].
- [95] Xiaolei Li and Arman Shafieloo, “Generalised Emergent Dark Energy Model: Confronting Λ and PEDE,” (2020), arXiv:2001.05103 [astro-ph.CO].
- [96] Giampaolo Benevento, Wayne Hu, and Marco Raveri, “Can Late Dark Energy Transitions Raise the Hubble constant?” (2020), arXiv:2002.11707 [astro-ph.CO].
- [97] <https://github.com/lkazantzi/pantheon-tomography>.